



## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

This report highlights all changes (additions, deletions, and removals) to the CVS Caremark® Advanced Control Formulary.

### ADDITIONS:

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Brand Agents:</b>			
<b>Abilify Maintena</b> (aripiprazole ext-rel) intramuscular extended-release suspension for injection	Central Nervous System/ Antipsychotics/ Atypicals	Abilify Maintena is indicated for: <ul style="list-style-type: none"> <li>• Treatment of schizophrenia in adults</li> <li>• Maintenance monotherapy treatment of bipolar I disorder in adults</li> </ul>	To provide a long-acting injectable antipsychotic option.
<b>Alecensa</b> (alectinib) oral capsule	Antineoplastic Agents/ Kinase Inhibitors	Alecensa is indicated for the treatment of patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer as detected by an FDA-approved test.	To provide an option for the treatment of ALK-positive non-small cell lung cancer.
<b>Alunbrig</b> (brigatinib) oral tablet, oral initiation pack	Antineoplastic Agents/ Kinase Inhibitors	Alunbrig is indicated for the treatment of adult patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer as detected by an FDA-approved test.	To provide an option for the treatment of ALK-positive non-small cell lung cancer.
<b>Annovera</b> (segesterone acetate- ethinyl estradiol) vaginal ring	Endocrine and Metabolic/ Contraceptives/ Vaginal	Annovera is indicated for use by females of reproductive potential to prevent pregnancy.	To provide an additional vaginal contraceptive option.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Bijuva</b> (estradiol-progesterone) oral capsule	Endocrine and Metabolic/ Menopausal Symptom Agents/ Oral	Bijuva is indicated in a woman with a uterus for the treatment of moderate to severe vasomotor symptoms due to menopause.	To provide an additional option for the treatment of moderate to severe vasomotor symptoms associated with menopause.
<b>Breztri Aerosphere</b> (budesonide-glycopyrrolate-formoterol) inhalation aerosol	Respiratory/ Anticholinergic / Beta Agonist / Steroid Inhalant Combinations	Breztri Aerosphere is indicated for the maintenance treatment of patients with chronic obstructive pulmonary disease.	To provide an additional triple therapy option for the maintenance treatment of chronic obstructive pulmonary disease.
<b>Clenpiq</b> (sodium picosulfate-magnesium oxide-citric acid) oral solution	Gastrointestinal/ Laxatives	Clenpiq is indicated for cleansing of the colon as a preparation for colonoscopy in adults and pediatric patients ages 9 years and older.	To provide an additional option for colon cleansing prior to a colonoscopy.
<b>Doptelet</b> (avatrombopag) oral tablet	Hematologic/ Thrombocytopenia Agents	Doptelet is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure</li> <li>• Thrombocytopenia in adult patients with chronic immune thrombocytopenia who have had an insufficient response to a previous treatment</li> </ul>	To provide an additional option for the treatment of thrombocytopenia.
<b>Duobrii</b> (halobetasol propionate-tazarotene) topical lotion	Topical/ Dermatology/ Antipsoriatics	Duobrii is indicated for the topical treatment of plaque psoriasis in adults.	To provide an additional topical option for the treatment of plaque psoriasis.
<b>Durolane</b> (sodium hyaluronate) intra-articular gel for injection	Analgesics/ Viscosupplements	Durolane is indicated for the treatment of pain in osteoarthritis of the knee in patients who have failed to respond adequately to	To provide an additional viscosupplement option for osteoarthritis.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		conservative non-pharmacological therapy or simple analgesics (e.g. acetaminophen).	
<b>Enstilar</b> (calcipotriene-betamethasone dipropionate) topical foam	Topical/ Dermatology/ Antipsoriatics	Enstilar is indicated for the topical treatment of plaque psoriasis in patients 12 years and older.	To provide an additional topical option for the treatment of plaque psoriasis.
<b>Erivedge</b> (vismodegib) oral capsule	Antineoplastic Agents/ Miscellaneous	Erivedge is indicated for the treatment of adults with metastatic basal cell carcinoma, or with locally advanced basal cell carcinoma that has recurred following surgery or who are not candidates for surgery and who are not candidates for radiation.	To provide an option for the treatment of metastatic basal cell carcinoma.
<b>Euflexxa</b> (sodium hyaluronate) intra-articular solution for injection	Analgesics/ Viscosupplements	Euflexxa is indicated for the treatment of pain in osteoarthritis of the knee in patients who have failed to respond adequately to conservative non-pharmacologic therapy and simple analgesics (e.g., acetaminophen).	To provide an additional viscosupplement option for osteoarthritis.
<b>Flarex</b> (fluorometholone acetate) ophthalmic suspension	Topical/ Ophthalmic/ Anti-Inflammatories/ Steroidal	Flarex is indicated for use in the treatment of steroid responsive inflammatory conditions of the palpebral and bulbar conjunctiva, cornea, and anterior segment of the eye.	To provide an additional ophthalmic anti-inflammatory option.
<b>Imvexxy</b> (estradiol) vaginal insert	Endocrine and Metabolic/ Menopausal	Imvexxy is indicated for the treatment of moderate to severe dyspareunia, a	To provide an additional vaginal option for the treatment of dyspareunia.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
	Symptom Agents/ Vaginal	symptom of vulvar and vaginal atrophy, due to menopause.	
<b>Inbrija</b> (levodopa) inhalation powder	Central Nervous System/ Antiparkinsonian Agents	Inbrija is indicated for the intermittent treatment of “off” episodes in patients with Parkinson's disease treated with carbidopa-levodopa.	To provide an option for the treatment of “off” episodes (return of Parkinson's symptoms) in those with Parkinson's disease.
<b>Kesimpta</b> (ofatumumab) subcutaneous solution for injection	Central Nervous System/ Multiple Sclerosis Agents	Kesimpta is indicated for the treatment of relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.	To provide an additional option for the treatment of relapsing forms of multiple sclerosis.
<b>Kevzara</b> (sarilumab) subcutaneous solution for injection	Immunologic Agents/ Autoimmune Agents/ Rheumatoid Arthritis	Kevzara is indicated for treatment of adult patients with moderately to severely active rheumatoid arthritis who have had an inadequate response or intolerance to one or more disease-modifying antirheumatic drugs (DMARDs).	To provide an additional option for the treatment of moderately to severely active rheumatoid arthritis.
<b>Latuda</b> (lurasidone) oral tablet	Central Nervous System/ Antipsychotics/ Atypicals	Latuda is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Schizophrenia in adults and adolescents (13 to 17 years)</li> <li>• Depressive episode associated with Bipolar I Disorder (bipolar depression) in adults and pediatric patients (10 to 17 years) as monotherapy</li> <li>• Depressive episode associated with Bipolar I Disorder (bipolar depression)</li> </ul>	To provide an additional oral antipsychotic option.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		in adults as adjunctive therapy with lithium or valproate	
<b>Nayzilam</b> (midazolam) nasal spray	Central Nervous System/ Anticonvulsants	Nayzilam is indicated for the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern in patients with epilepsy 12 years of age and older.	To provide an additional option for the treatment of acute, intermittent seizures.
<b>Nexletol</b> (bempedoic acid) oral tablet	Cardiovascular/ Antilipemics/ ACL Inhibitors / Combinations	Nexletol is indicated as an adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or established atherosclerotic cardiovascular disease who require additional lowering of low-density lipoprotein cholesterol (LDL-C).	To provide an adjunctive option for the treatment of atherosclerotic cardiovascular disease and familial hypercholesterolemia.
<b>Nexlizet</b> (bempedoic acid-ezetimibe) oral tablet	Cardiovascular/ Antilipemics/ ACL Inhibitors / Combinations	Nexlizet is indicated as an adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or established atherosclerotic cardiovascular disease who require additional lowering of low-density lipoprotein cholesterol (LDL-C).	To provide an adjunctive option for the treatment of atherosclerotic cardiovascular disease and familial hypercholesterolemia.
<b>Ninlaro</b> (ixazomib) oral capsule	Antineoplastic Agents/ Multiple Myeloma/ Proteasome Inhibitors	Ninlaro is indicated in combination with lenalidomide and dexamethasone for the treatment of patients with multiple myeloma	To provide an additional option for the treatment of multiple myeloma.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		who have received at least one prior therapy.	
<b>Norditropin</b> (somatropin) subcutaneous solution for injection	Endocrine and Metabolic/ Human Growth Hormones	Norditropin is indicated for: <ul style="list-style-type: none"> <li>• Pediatric: Treatment of pediatric patients with growth failure due to inadequate secretion of endogenous growth hormone (GH), short stature associated with Noonan syndrome, short stature associated with Turner syndrome, short stature born small for gestational age with no catch-up growth by age 2 to 4 years, Idiopathic Short Stature, and growth failure due to Prader-Willi Syndrome</li> <li>• Adult: Replacement of endogenous GH in adults with growth hormone deficiency</li> </ul>	To provide an option for the treatment of growth hormone deficiency.
<b>Ocrevus</b> (ocrelizumab) intravenous solution for injection	Central Nervous System/ Multiple Sclerosis Agents	Ocrevus is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults</li> <li>• Primary progressive multiple sclerosis, in adults</li> </ul>	To provide an option for the treatment of primary progressive multiple sclerosis and an additional option for the treatment of relapsing forms of multiple sclerosis.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Omnipod Insulin Infusion Pump</b> insulin infusion disposable pump	Endocrine and Metabolic/ Antidiabetics/ Supplies	Omnipod is used to allow continuous subcutaneous basal insulin infusion and on-demand bolus dosing in those with insulin-dependent diabetes.	To provide an additional continuous/basal and on-demand/bolus insulin delivery option in insulin-dependent diabetes.
<b>Omnipod DASH Insulin Infusion Pump</b> insulin management system	Endocrine and Metabolic/ Antidiabetics/ Supplies	Omnipod DASH is used to allow continuous subcutaneous basal insulin infusion and on-demand bolus dosing in those with insulin-dependent diabetes.	To provide an additional continuous/basal and on-demand/bolus insulin delivery option in insulin-dependent diabetes.
<b>OneTouch Lancets</b> lancets	Endocrine and Metabolic/ Antidiabetics/ Supplies	OneTouch lancets are supplies that aid in the testing of blood glucose levels in those who have diabetes.	To provide a preferred lancet option for testing blood glucose levels.
<b>OneTouch Lancing Devices</b> lancing devices	Endocrine and Metabolic/ Antidiabetics/ Supplies	OneTouch lancing devices are supplies that aid in the testing of blood glucose levels in those who have diabetes.	To provide a preferred lancing device option for testing blood glucose levels.
<b>OneTouch Ultra, OneTouch Verio Strips and Kits</b> blood glucose test strips, monitoring kits	Endocrine and Metabolic/ Antidiabetics/ Supplies	OneTouch Ultra and OneTouch Verio strips and kits are used to test and monitor blood glucose levels in those who have diabetes.	To provide an option for testing and monitoring blood glucose levels.
<b>Oracea</b> (doxycycline monohydrate delayed-rel) oral delayed-release capsule	Topical/ Dermatology/ Rosacea	Oracea is indicated for the treatment of only inflammatory lesions (papules and pustules) of rosacea in adult patients.	To provide an oral option for the treatment of rosacea.
<b>Perjeta</b> (pertuzumab) intravenous solution for injection	Antineoplastic Agents/ Miscellaneous	Perjeta is indicated for: <ul style="list-style-type: none"> <li>• Use in combination with trastuzumab and docetaxel for treatment of patients</li> </ul>	To provide an additional option for the treatment of human epidermal growth factor receptor 2 (HER2)-positive breast cancer.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<p>with HER2-positive metastatic breast cancer who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease</p> <ul style="list-style-type: none"> <li>• Use in combination with trastuzumab and chemotherapy as:               <ul style="list-style-type: none"> <li>○ Neoadjuvant treatment of patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) as part of a complete treatment regimen for early breast cancer</li> <li>○ Adjuvant treatment of patients with HER2-positive early breast cancer at high risk of recurrence</li> </ul> </li> </ul>	
<b>Perseris</b> (risperidone ext-rel) subcutaneous extended-release suspension for injection	Central Nervous System/ Antipsychotics/ Atypicals	Perseris is indicated for the treatment of schizophrenia in adults.	To provide a long-acting injectable antipsychotic option.
<b>Phesgo</b> (pertuzumab-trastuzumab-hyaluronidase-zzxf) subcutaneous solution for injection	Antineoplastic Agents/ Miscellaneous	Phesgo is indicated for: <ul style="list-style-type: none"> <li>• Use in combination with chemotherapy as:               <ul style="list-style-type: none"> <li>○ Neoadjuvant treatment of patients with HER2-positive, locally</li> </ul> </li> </ul>	To provide an additional option for the treatment of human epidermal growth factor receptor 2 (HER2)-positive breast cancer.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.



Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<p>advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) as part of a complete treatment regimen for early breast cancer</p> <ul style="list-style-type: none"> <li>○ Adjuvant treatment of patients with HER2-positive early breast cancer at high risk of recurrence</li> <li>● Use in combination with docetaxel for treatment of patients with HER2-positive metastatic breast cancer who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease</li> </ul>	
<b>Sancuso</b> (granisetron) transdermal patch	Gastrointestinal/ Antiemetics	Sancuso is indicated for the prevention of nausea and vomiting in adults receiving moderately and/or highly emetogenic chemotherapy for up to 5 consecutive days.	To provide an additional option for the prevention of chemotherapy-induced nausea and vomiting.
<b>Saphris</b> (asenapine) sublingual tablet	Central Nervous System/ Antipsychotics/ Atypicals	<p>Saphris is indicated for:</p> <ul style="list-style-type: none"> <li>● Schizophrenia in adults</li> <li>● Bipolar I disorder               <ul style="list-style-type: none"> <li>○ Acute monotherapy treatment of manic or mixed episodes, in adults and pediatric patients 10 to 17 years of age</li> </ul> </li> </ul>	To provide an additional oral antipsychotic option.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<ul style="list-style-type: none"> <li>○ Adjunctive treatment to lithium or valproate in adults</li> <li>○ Maintenance monotherapy treatment in adults</li> </ul>	
<b>Simbrinza</b> (brinzolamide-brimonidine) ophthalmic suspension	Topical/ Ophthalmic/ Carbonic Anhydrase Inhibitor / Sympathomimetic Combinations	Simbrinza is indicated for the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension.	To provide an additional option for the reduction of elevated intraocular pressure.
<b>Taclonex</b> (calcipotriene-betamethasone dipropionate) topical ointment	Topical/ Dermatology/ Antipsoriatics	Taclonex ointment is indicated for the topical treatment of plaque psoriasis in patients 12 years of age and older.	To provide an additional topical option for the treatment of plaque psoriasis.
<b>Taclonex</b> (calcipotriene-betamethasone dipropionate) topical suspension	Topical/ Dermatology/ Antipsoriatics	Taclonex suspension is indicated for the topical treatment of plaque psoriasis of the scalp and body in patients 12 years and older.	To provide an additional topical option for the treatment of plaque psoriasis.
<b>Toujeo</b> (insulin glargine) subcutaneous solution for injection	Endocrine and Metabolic/ Antidiabetics/ Insulins	Toujeo is indicated to improve glycemic control in adults and pediatric patients 6 years and older with diabetes mellitus.	To provide an additional long-acting insulin option for the management of diabetes mellitus.
<b>Valtoco</b> (diazepam) nasal spray	Central Nervous System/ Anticonvulsants	Valtoco is indicated for the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are	To provide an additional option for the treatment of acute, intermittent seizures.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		distinct from a patient's usual seizure pattern in patients with epilepsy 6 years of age and older.	
<b>Velcade</b> (bortezomib) intravenous / subcutaneous solution for injection	Antineoplastic Agents/ Multiple Myeloma/ Proteasome Inhibitors	Velcade is indicated for: <ul style="list-style-type: none"> <li>• Treatment of adult patients with multiple myeloma</li> <li>• Treatment of adult patients with mantle cell lymphoma</li> </ul>	To provide an additional option for the treatment of multiple myeloma and mantle cell lymphoma.
<b>Xcopri</b> (cenobamate) oral tablet, oral titration pack, oral maintenance pack	Central Nervous System/ Anticonvulsants	Xcopri is indicated for the treatment of partial-onset seizures in adult patients.	To provide an additional option for the treatment of partial-onset seizures in adults.
<b>Xospata</b> (gilteritinib fumarate) oral tablet	Antineoplastic Agents/ Kinase Inhibitors	Xospata is indicated for the treatment of adult patients who have relapsed or refractory acute myeloid leukemia with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test.	To provide an option for the treatment of relapsed or refractory acute myeloid leukemia with a specific type of genetic mutation.
<b>Zeposia</b> (ozanimod) oral capsule, oral starter pack	Central Nervous System/ Multiple Sclerosis Agents	Zeposia is indicated for the treatment of relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.	To provide an additional option for the treatment of relapsing forms of multiple sclerosis.
<b>Ziextenzo</b> (pegfilgrastim-bmez) subcutaneous solution for injection	Hematologic/ Hematopoietic Growth Factors	Ziextenzo is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs	To provide a long-acting colony-stimulating factor option for those who are receiving myelosuppressive anti-cancer therapy.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		associated with a clinically significant incidence of febrile neutropenia.	
<b>Generic Agents:</b>			
<b>aprepitant</b> oral capsule	Gastrointestinal/ Antiemetics	Aprepitant is indicated: <ul style="list-style-type: none"> <li>• In combination with other antiemetic agents, in patients 12 years of age and older for prevention of:               <ul style="list-style-type: none"> <li>○ Acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy including high-dose cisplatin</li> <li>○ Nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy</li> </ul> </li> <li>• For prevention of postoperative nausea and vomiting in adults</li> </ul>	To provide an additional generic option for the prevention of chemotherapy-induced nausea and vomiting and postoperative nausea and vomiting.
<b>ciprofloxacin-dexamethasone</b> otic suspension	Topical/ Otic/ Anti- Infective / Anti- Inflammatory Combinations	Ciprofloxacin-dexamethasone is indicated for the treatment of infections caused by susceptible isolates of the designated microorganisms in the specific conditions listed below: <ul style="list-style-type: none"> <li>• Acute Otitis Media (AOM) in pediatric patients (age 6 months and older) with tympanostomy tubes due to <i>Staphylococcus aureus</i>, <i>Streptococcus</i></li> </ul>	To provide an additional generic otic anti-infective and anti-inflammatory option for the treatment of ear infections.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<p><i>pneumoniae, Haemophilus influenzae, Moraxella catarrhalis, and Pseudomonas aeruginosa</i></p> <ul style="list-style-type: none"> <li>Acute Otitis Externa (AOE) in pediatric (age 6 months and older), adult and elderly patients due to <i>Staphylococcus aureus</i> and <i>Pseudomonas aeruginosa</i></li> </ul>	
<p><b>dimethyl fumarate delayed-rel</b> oral delayed-release capsule</p>	<p>Central Nervous System/ Multiple Sclerosis Agents</p>	<p>Dimethyl fumarate is indicated for the treatment of relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.</p>	<p>To provide an additional generic option for the treatment of relapsing forms of multiple sclerosis.</p>
<p><b>isosorbide mononitrate</b> oral tablet</p>	<p>Cardiovascular/ Nitrates/ Oral</p>	<p>Isosorbide mononitrate is indicated for the prevention and treatment of angina pectoris due to coronary artery disease.</p>	<p>To provide an additional oral nitrate option.</p>
<p><b>pyrimethamine</b> oral tablet</p>	<p>Anti-Infectives/ Miscellaneous</p>	<p>Pyrimethamine is indicated for the treatment of toxoplasmosis when used conjointly with a sulfonamide, since synergism exists with this combination.</p>	<p>To provide a generic option for the treatment of toxoplasmosis.</p>

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

**DELETIONS:**

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Brand Agents:</b>			
<b>Accu-Chek Lancets</b> lancets	Endocrine and Metabolic/ Antidiabetics/ Supplies	Accu-Chek lancets are supplies that aid in the testing of blood glucose levels in those who have diabetes.	Availability of an additional lancet option for testing blood glucose levels.  The preferred option on the Advanced Control Formulary is OneTouch lancets.
<b>Accu-Chek Lancing Devices</b> lancing devices	Endocrine and Metabolic/ Antidiabetics/ Supplies	Accu-Chek lancing devices are supplies that aid in the testing of blood glucose levels in those who have diabetes.	Availability of an additional lancing device option for testing blood glucose levels.  The preferred option on the Advanced Control Formulary is OneTouch Lancing Devices.
<b>Aristada</b> (aripiprazole lauroxil ext-rel) intramuscular extended-release suspension for injection	Central Nervous System/ Antipsychotics/ Atypicals	Aristada is indicated for the treatment of schizophrenia in adults.	Availability of additional long-acting injectable antipsychotic options.  Preferred options on the Advanced Control Formulary include Abilify Maintena (aripiprazole ext-rel injection) and Perseris (risperidone ext-rel injection).
<b>Aristada Initio</b> (aripiprazole lauroxil ext-rel) intramuscular extended-release suspension for injection	Central Nervous System/ Antipsychotics/ Atypicals	Aristada Initio, in combination with oral aripiprazole, is indicated for the initiation of Aristada when used for the treatment of schizophrenia in adults.	Availability of additional long-acting injectable antipsychotic options.  Preferred options on the Advanced Control Formulary include Abilify Maintena (aripiprazole ext-rel injection) and Perseris (risperidone ext-rel injection).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

**REMOVALS:**

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Brand Agents:</b>			
<b>Accu-Chek Aviva Plus, Accu-Chek Compact Plus, Accu-Chek Guide, Accu-Chek SmartView Strips and Kits</b> blood glucose test strips, monitoring kits	Endocrine and Metabolic/ Antidiabetics/ Supplies	Accu-Chek strips and kits are used to test and monitor blood glucose levels in those who have diabetes.	Availability of additional options for testing and monitoring blood glucose levels.  Preferred options on the Advanced Control Formulary include OneTouch Ultra strips and kits and OneTouch Verio strips and kits.
<b>Adzenys ER</b> (amphetamine ext-rel) oral extended-release suspension	Central Nervous System/ Attention Deficit Hyperactivity Disorder	Adzenys ER is indicated for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 6 years and older.	Availability of additional options for the treatment of Attention Deficit Hyperactivity Disorder.  Preferred options on the Advanced Control Formulary include amphetamine-dextroamphetamine mixed salts ext-rel (excluding certain NDCs), dexamethylphenidate ext-rel, dextroamphetamine ext-rel, methylphenidate ext-rel (excluding certain NDCs), Mydayis (amphetamine-dextroamphetamine mixed salts ext-rel), and Vyvanse (lisdexamfetamine).
<b>Akynzeo</b> (netupitant-palonosetron) oral capsule	Gastrointestinal/ Antiemetics	Akynzeo capsule is indicated in combination with dexamethasone in adults for the prevention of acute and delayed nausea and vomiting associated with initial and repeat courses of cancer chemotherapy, including, but not limited to, highly emetogenic chemotherapy.	Availability of additional options for the prevention of chemotherapy-induced nausea and vomiting.  Preferred options on the Advanced Control Formulary include aprepitant WITH granisetron, ondansetron, or Sancuso (granisetron).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Akynzeo</b> (fosnetupitant-palonosetron) intravenous solution for injection	Gastrointestinal/ Antiemetics	Akynzeo injection is indicated in combination with dexamethasone in adults for the prevention of acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy.	Availability of additional options for the prevention of chemotherapy-induced nausea and vomiting.  Preferred options on the Advanced Control Formulary include aprepitant WITH granisetron, ondansetron, or Sancuso (granisetron).
<b>Amitiza</b> (lubiprostone) oral capsule	Gastrointestinal/ Irritable Bowel Syndrome	Amitiza is indicated for the treatment of: <ul style="list-style-type: none"> <li>• Chronic idiopathic constipation (CIC) in adults</li> <li>• Opioid-induced constipation (OIC) in adult patients with chronic, non-cancer pain, including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation</li> <li>• Irritable bowel syndrome with constipation (IBS-C) in women ≥ 18 years old</li> </ul>	Availability of additional options for the treatment of chronic idiopathic constipation, opioid-induced constipation, and irritable bowel syndrome with constipation.  Preferred options on the Advanced Control Formulary include Linzess (linaclotide), Movantik (naloxegol), and Symproic (naldemedine).
<b>Anzemet</b> (dolasetron mesylate) oral tablet	Gastrointestinal/ Antiemetics	Anzemet is indicated for the prevention of nausea and vomiting associated with moderately emetogenic cancer chemotherapy, including initial and repeat courses in adults and children 2 years and older.	Availability of additional options for the prevention of chemotherapy-induced nausea and vomiting.  Preferred options on the Advanced Control Formulary include granisetron, ondansetron, and Sancuso (granisetron).
<b>Apokyn</b> (apomorphine) subcutaneous solution for injection	Central Nervous System/ Antiparkinsonian Agents	Apokyn is indicated for the acute, intermittent treatment of hypomobility, "off" episodes ("end-of-dose wearing off" and	Availability of an additional option for the treatment of "off" episodes (return of Parkinson's symptoms) in those with advanced Parkinson's disease.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.



## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		unpredictable "on/off" episodes) associated with advanced Parkinson's disease.	The preferred option the Advanced Control Formulary is Inbrija (levodopa inhalation powder).
<b>Aralast NP</b> (alpha-1 proteinase inhibitor) intravenous solution for injection	Respiratory/ Alpha-1 Antitrypsin Deficiency Agents	Aralast NP is indicated for chronic augmentation therapy in adults with clinically evident emphysema due to severe congenital deficiency of alpha <sub>1</sub> -antitrypsin inhibitor.	Availability of an additional option for the treatment of emphysema due to an inherited disorder known as alpha <sub>1</sub> -antitrypsin deficiency.  The preferred option on the Advanced Control Formulary is Prolastin-C (alpha-1 proteinase inhibitor).
<b>Azesco</b> (multivitamin with iron) oral tablet	Nutritional/Supplements/ Vitamins and Minerals/ Prenatal Vitamins	Azesco is used: <ul style="list-style-type: none"> <li>• For the clinical dietary management of suboptimal nutritional status in patients where advanced folate supplementation is required and nutritional supplementation in physiologically stressful conditions for maintenance of good health is needed</li> <li>• Throughout pregnancy, during the postnatal period for both lactating and non-lactating mothers, and throughout the childbearing years</li> <li>• For improving the nutritional status of women prior to conception</li> </ul>	Availability of additional prenatal vitamin options.  Preferred options on the Advanced Control formulary include prenatal vitamins and Citranatal (prenatal vitamins with folic acid).
<b>Azopt</b> (brinzolamide) ophthalmic suspension	Topical/ Ophthalmic/ Carbonic Anhydrase Inhibitors	Azopt is indicated for the treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma.	Availability of a generic ophthalmic carbonic anhydrase inhibitor option.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
			The preferred option on the Advanced Control Formulary is dorzolamide.
<b>Besivance</b> (besifloxacin) ophthalmic suspension	Topical/ Ophthalmic/ Anti-Infectives	Besivance is indicated for the treatment of bacterial conjunctivitis caused by susceptible isolates of the following bacteria: <i>Aerococcus viridans</i> , CDC coryneform group G, <i>Corynebacterium pseudodiphtheriticum</i> , <i>Corynebacterium striatum</i> , <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i> , <i>Moraxella lacunata</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Staphylococcus hominis</i> , <i>Staphylococcus lugdunensis</i> , <i>Staphylococcus warneri</i> , <i>Streptococcus mitis group</i> , <i>Streptococcus oralis</i> , <i>Streptococcus pneumoniae</i> , <i>Streptococcus salivarius</i> .	Availability of generic ophthalmic anti-infective options.  Preferred options on the Advanced Control Formulary include ciprofloxacin, erythromycin, gentamicin, levofloxacin, moxifloxacin, ofloxacin, sulfacetamide, and tobramycin.
<b>Betoptic S</b> (betaxolol) ophthalmic suspension	Topical/ Ophthalmic/ Beta-Blockers/ Selective	Betoptic S is indicated for the treatment of elevated intraocular pressure (IOP) in patients with chronic open-angle glaucoma or ocular hypertension.	Availability of additional ophthalmic beta-blocker options.  Preferred options on the Advanced Control Formulary include timolol maleate solution and Betimol (timolol hemihydrate).
<b>Bevespi Aerosphere</b> (glycopyrrolate-formoterol) inhalation aerosol	Respiratory/ Anticholinergic / Beta Agonist Combinations/ Long Acting	Bevespi Aerosphere is indicated for the maintenance treatment of patients with chronic obstructive pulmonary disease (COPD).	Availability of additional dual therapy options for the maintenance treatment of chronic obstructive pulmonary disease.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
			Preferred options on the Advanced Control Formulary include Anoro Ellipta (umeclidinium-vilanterol) and Stiolto Respimat (tiotropium-olodaterol).
<b>Bortezomib</b> (bortezomib) intravenous solution for injection	Antineoplastic Agents/ Multiple Myeloma/ Proteasome Inhibitors	Bortezomib is indicated for: <ul style="list-style-type: none"> <li>• Treatment of patients with multiple myeloma</li> <li>• Treatment of patients with mantle cell lymphoma who have received at least 1 prior therapy</li> </ul>	Availability of additional options for the treatment of multiple myeloma and mantle cell lymphoma.  Preferred options on the Advanced Control Formulary include Ninlaro (ixazomib) and Velcade (bortezomib).
<b>Briviact</b> (brivaracetam) oral tablet, oral solution	Central Nervous System/ Anticonvulsants	Briviact oral tablet and oral solution are indicated for the treatment of partial-onset seizures in patients 4 years of age and older.	Availability of additional anticonvulsant options.  Preferred options on the Advanced Control Formulary include carbamazepine, carbamazepine ext-rel, divalproex sodium, divalproex sodium ext-rel, gabapentin, lamotrigine, lamotrigine ext-rel, levetiracetam, levetiracetam ext-rel, oxcarbazepine, phenobarbital, phenytoin, phenytoin sodium extended, primidone, tiagabine, topiramate, valproic acid, zonisamide, Oxtellar XR (oxcarbazepine ext-rel), Trokendi XR (topiramate ext-rel), Vimpat (lacosamide), and Xcopri (cenobamate).
<b>Briviact</b> (brivaracetam) intravenous solution for injection	Central Nervous System/ Anticonvulsants	Briviact injection is indicated for the treatment of partial-onset seizures only in adult patients (16 years of age and older).	Availability of additional anticonvulsant options.  Preferred options on the Advanced Control Formulary include carbamazepine, carbamazepine ext-rel, divalproex sodium, divalproex sodium ext-rel, gabapentin, lamotrigine, lamotrigine ext-rel,

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
			levetiracetam, levetiracetam ext-rel, oxcarbazepine, phenobarbital, phenytoin, phenytoin sodium extended, primidone, tiagabine, topiramate, valproic acid, zonisamide, Oxtellar XR (oxcarbazepine ext-rel), Trokendi XR (topiramate ext-rel), Vimpat (lacosamide), and Xcopri (cenobamate).
<b>Ciloxan</b> (ciprofloxacin) ophthalmic ointment	Topical/ Ophthalmic/ Anti-Infectives	Ciloxan ointment is indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the microorganisms listed below: <ul style="list-style-type: none"> <li>• Gram-Positive: <i>Staphylococcus aureus</i>, <i>Staphylococcus epidermidis</i>, <i>Streptococcus pneumoniae</i>, <i>Streptococcus</i> (Viridans Group)</li> <li>• Gram-Negative: <i>Haemophilus influenzae</i></li> </ul>	Availability of generic ophthalmic anti-infective options.  Preferred options on the Advanced Control Formulary include ciprofloxacin, erythromycin, gentamicin, levofloxacin, moxifloxacin, ofloxacin, sulfacetamide, and tobramycin.
<b>Ciloxan</b> (ciprofloxacin) ophthalmic solution	Topical/ Ophthalmic/ Anti-Infectives	Ciloxan solution is indicated for the treatment of infections caused by susceptible strains of the designated microorganisms in the conditions listed below: <ul style="list-style-type: none"> <li>• Corneal Ulcers: <i>Pseudomonas aeruginosa</i>, <i>Serratia marcescens</i>, <i>Staphylococcus aureus</i>, <i>Staphylococcus epidermidis</i>, <i>Streptococcus pneumoniae</i>, <i>Streptococcus</i> (Viridans Group)</li> </ul>	Availability of generic ophthalmic anti-infective options.  Preferred options on the Advanced Control Formulary include ciprofloxacin, erythromycin, gentamicin, levofloxacin, moxifloxacin, ofloxacin, sulfacetamide, and tobramycin.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<ul style="list-style-type: none"> <li>Conjunctivitis: <i>Haemophilus influenzae</i>, <i>Staphylococcus aureus</i>, <i>Staphylococcus epidermidis</i>, <i>Streptococcus pneumoniae</i></li> </ul>	
<b>Cipro HC Otic</b> (ciprofloxacin-hydrocortisone) otic suspension	Topical/ Otic/ Anti-Infective / Anti-Inflammatory Combinations	Cipro HC Otic is indicated for the treatment of acute otitis externa in adult and pediatric patients, one year and older, due to susceptible strains of <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , and <i>Proteus mirabilis</i> .	Availability of generic otic anti-infective and/or anti-inflammatory options.  Preferred options on the Advanced Control Formulary include ciprofloxacin-dexamethasone and ofloxacin otic.
<b>Ciprodex</b> (ciprofloxacin-dexamethasone) otic suspension	Topical/ Otic/ Anti-Infective / Anti-Inflammatory Combinations	Ciprodex is indicated for the treatment of infections caused by susceptible isolates of the designated microorganisms in the specific conditions listed below: <ul style="list-style-type: none"> <li>Acute Otitis Media (AOM) in pediatric patients (age 6 months and older) with tympanostomy tubes due to <i>Staphylococcus aureus</i>, <i>Streptococcus pneumoniae</i>, <i>Haemophilus influenzae</i>, <i>Moraxella catarrhalis</i>, and <i>Pseudomonas aeruginosa</i></li> <li>Acute Otitis Externa (AOE) in pediatric (age 6 months and older), adult and elderly patients due to <i>Staphylococcus aureus</i> and <i>Pseudomonas aeruginosa</i></li> </ul>	Availability of generic otic anti-infective and/or anti-inflammatory options.  Preferred options on the Advanced Control Formulary include ciprofloxacin-dexamethasone and ofloxacin otic.
<b>Daraprim</b> (pyrimethamine)	Anti-Infectives/ Miscellaneous	Daraprim is indicated for the treatment of toxoplasmosis when used conjointly with a	Availability of a generic option for the treatment of toxoplasmosis.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
oral tablet		sulfonamide, since synergism exists with this combination.	The preferred option on the Advanced Control Formulary is pyrimethamine.
<b>Daytrana</b> (methylphenidate) transdermal patch	Central Nervous System/ Attention Deficit Hyperactivity Disorder	Daytrana is indicated for the treatment of Attention Deficit Hyperactivity Disorder (ADHD).	Availability of additional options for the treatment of Attention Deficit Hyperactivity Disorder.  Preferred options on the Advanced Control Formulary include amphetamine-dextroamphetamine mixed salts ext-rel (excluding certain NDCs), dexamethylphenidate ext-rel, dextroamphetamine ext-rel, methylphenidate ext-rel (excluding certain NDCs), Mydayis (amphetamine-dextroamphetamine mixed salts ext-rel), and Vyvanse (lisdexamfetamine).
<b>Depo-subQ Provera 104</b> (medroxyprogesterone acetate) subcutaneous suspension for injection	Endocrine and Metabolic/ Contraceptives/ Injectable	Depo-subQ Provera is indicated in females of reproductive age for: <ul style="list-style-type: none"> <li>• Prevention of pregnancy</li> <li>• Management of endometriosis-associated pain</li> </ul>	Availability of a generic injectable option for contraception and the management of endometriosis-associated pain.  The preferred option on the Advanced Control Formulary is medroxyprogesterone acetate 150 mg/mL.
<b>Differin Lotion</b> (adapalene) topical lotion	Topical/ Dermatology/ Acne/ Topical	Differin Lotion is indicated for the topical treatment of acne vulgaris in patients 12 years and older.	Availability of additional options for the topical treatment of acne.  Preferred options on the Advanced Control Formulary include adapalene, benzoyl peroxide, clindamycin gel (except NDC 68682046275), clindamycin solution, clindamycin-benzoyl peroxide, erythromycin solution, erythromycin-benzoyl peroxide, tretinoin, Epiduo

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
			(adapalene-benzoyl peroxide), and Onexton (clindamycin-benzoyl peroxide).
<b>Duavee</b> (conjugated estrogens-bazedoxifene) oral tablet	Endocrine and Metabolic/ Menopausal Symptom Agents/ Oral	Duavee is indicated for treatment of the following conditions in women with a uterus: <ul style="list-style-type: none"> <li>• Treatment of moderate to severe vasomotor symptoms associated with menopause</li> <li>• Prevention of postmenopausal osteoporosis</li> </ul>	Availability of additional options for the treatment of vasomotor symptoms associated with menopause and prevention of postmenopausal osteoporosis.  Preferred options on the Advanced Control Formulary include estradiol-norethindrone, raloxifene, and Bijuva (estradiol-progesterone).
<b>Emend</b> (aprepitant) oral capsule	Gastrointestinal/ Antiemetics	Emend capsule is indicated in combination with other antiemetic agents, in patients 12 years of age and older for prevention of: <ul style="list-style-type: none"> <li>• Acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy including high-dose cisplatin</li> <li>• Nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy</li> </ul>	Availability of a generic option for the prevention of chemotherapy-induced nausea and vomiting.  The preferred option on the Advanced Control Formulary is aprepitant.
<b>Emend</b> (aprepitant) powder for oral suspension	Gastrointestinal/ Antiemetics	Emend suspension is indicated in combination with other antiemetic agents, in patients 6 months of age and older for prevention of: <ul style="list-style-type: none"> <li>• Acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer</li> </ul>	Availability of a generic option for the prevention of chemotherapy-induced nausea and vomiting.  The preferred option on the Advanced Control Formulary is aprepitant.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		chemotherapy including high-dose cisplatin <ul style="list-style-type: none"> <li>Nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy</li> </ul>	
<b>Emend</b> (fosaprepitant) intravenous solution for injection	Gastrointestinal/ Antiemetics	Emend injection is indicated in adults and pediatric patients 6 months of age and older, in combination with other antiemetic agents, for the prevention of: <ul style="list-style-type: none"> <li>Acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy including high-dose cisplatin</li> <li>Delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy</li> </ul>	Availability of a generic option for the prevention of chemotherapy-induced nausea and vomiting.  The preferred option on the Advanced Control Formulary is aprepitant.
<b>Fycompa</b> (perampanel) oral tablet, oral suspension	Central Nervous System/ Anticonvulsants	Fycompa is indicated for: <ul style="list-style-type: none"> <li>Treatment of partial-onset seizures with or without secondarily generalized seizures in patients with epilepsy 4 years of age and older</li> <li>Adjunctive therapy in the treatment of primary generalized tonic-clonic seizures in patients with epilepsy 12 years of age and older</li> </ul>	Availability of additional anticonvulsant options.  Preferred options on the Advanced Control Formulary include carbamazepine, carbamazepine ext-rel, divalproex sodium, divalproex sodium ext-rel, gabapentin, lamotrigine, lamotrigine ext-rel, levetiracetam, levetiracetam ext-rel, oxcarbazepine, phenobarbital, phenytoin, phenytoin sodium extended, primidone, tiagabine, topiramate, valproic acid,

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.



## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
			zonisamide, Oxtellar XR (oxcarbazepine ext-rel), Trokendi XR (topiramate ext-rel), Vimpat (lacosamide), and Xcopri (cenobamate).
<b>Gel-One</b> (sodium hyaluronate) intra-articular gel for injection	Analgesics/ Viscosupplements	Gel-One is indicated for the treatment of pain in osteoarthritis of the knee in patients who have failed to respond adequately to non-pharmacologic therapy, non-steroidal anti-inflammatory drugs (NSAIDs) or simple analgesics (e.g., acetaminophen).	Availability of additional viscosupplement options for osteoarthritis.  Preferred options on the Advanced Control Formulary include Durolane (sodium hyaluronate), Euflexxa (sodium hyaluronate), Gelsyn-3 (sodium hyaluronate), and Supartz FX (sodium hyaluronate).
<b>Glassia</b> (alpha-1 proteinase inhibitor) intravenous solution for injection	Respiratory/ Alpha-1 Antitrypsin Deficiency Agents	Glassia is indicated for chronic augmentation and maintenance therapy in adults with clinically evident emphysema due to severe hereditary deficiency of alpha <sub>1</sub> -antitrypsin inhibitor.	Availability of an additional option for the treatment of emphysema due to an inherited disorder known as alpha <sub>1</sub> -antitrypsin deficiency.  The preferred option on the Advanced Control Formulary is Prolastin-C (alpha-1 proteinase inhibitor).
<b>GoLYTELY</b> (peg 3350-electrolytes) powder for oral solution	Gastrointestinal/ Laxatives	GoLYTELY is indicated for cleansing of the colon in preparation for colonoscopy and barium enema X-ray examination in adults.	Availability of additional options for colon cleansing prior to a colonoscopy.  Preferred options on the Advanced Control Formulary include peg 3350-electrolytes and Clenpiq (sodium picosulfate-magnesium oxide-citric acid).
<b>Humatrope</b> (somatropin) subcutaneous solution for injection	Endocrine and Metabolic/ Human Growth Hormones	Humatrope is indicated for: <ul style="list-style-type: none"> <li>• Pediatric: growth failure due to inadequate secretion of endogenous growth hormone (GH); short stature associated with Turner syndrome;</li> </ul>	Availability of an additional option for the treatment of growth hormone deficiency.  The preferred option on the Advanced Control Formulary is Norditropin (somatropin).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		Idiopathic Short Stature, height standard deviation score <-2.25, and associated with growth rates unlikely to permit attainment of adult height in the normal range; short stature or growth failure in short stature homeobox-containing gene deficiency; short stature born small for gestational age with no catch-up growth by 2 years to 4 years of age <ul style="list-style-type: none"> <li>Adult: replacement of endogenous GH in adults with GH deficiency</li> </ul>	
<b>Incruse Ellipta</b> (umeclidinium bromide) oral inhalation powder	Respiratory/ Anticholinergics	Incruse Ellipta is indicated for the maintenance treatment of patients with chronic obstructive pulmonary disease (COPD).	Availability of additional long-acting anticholinergic options for the maintenance treatment of chronic obstructive pulmonary disease.  Preferred options on the Advanced Control Formulary include Spiriva (tiotropium) and Yupelri (revefenacin inhalation solution).
<b>Intrarosa</b> (prasterone) vaginal insert	Endocrine and Metabolic/ Menopausal Symptom Agents/ Vaginal	Intrarosa is indicated for the treatment of moderate to severe dyspareunia, a symptom of vulvar and vaginal atrophy, due to menopause.	Availability of additional vaginal options for the treatment of dyspareunia.  Preferred options on the Advanced Control Formulary include estradiol and Imvexxy (estradiol).
<b>Invega Sustenna</b> (paliperidone palmitate ext-rel)	Central Nervous System/ Antipsychotics/ Atypicals	Invega Sustenna is indicated for: <ul style="list-style-type: none"> <li>Treatment of schizophrenia in adults</li> </ul>	Availability of additional long-acting injectable antipsychotic options.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
intramuscular extended-release suspension for injection		<ul style="list-style-type: none"> <li>Treatment of schizoaffective disorder in adults as monotherapy and as an adjunct to mood stabilizers or antidepressants</li> </ul>	Preferred options on the Advanced Control Formulary include Abilify Maintena (aripiprazole ext-rel injection) and Perseris (risperidone ext-rel injection).
<b>Kyprolis</b> (carfilzomib) intravenous solution for injection	Antineoplastic Agents/ Multiple Myeloma/ Proteasome Inhibitors	Kyprolis is indicated: <ul style="list-style-type: none"> <li>In combination with dexamethasone or with lenalidomide plus dexamethasone for the treatment of patients with relapsed or refractory multiple myeloma who have received one to three lines of therapy</li> <li>As a single agent for the treatment of patients with relapsed or refractory multiple myeloma who have received one or more lines of therapy</li> </ul>	Availability of additional options for the treatment of relapsed or refractory multiple myeloma.  Preferred options on the Advanced Control Formulary include Ninlaro (ixazomib) and Velcade (bortezomib).
<b>Lacrisert</b> (hydroxypropyl cellulose) ophthalmic insert	Topical/ Ophthalmic/ Artificial Tears	Lacrisert is indicated in: <ul style="list-style-type: none"> <li>Moderate to severe dry eye syndromes, including keratoconjunctivitis sicca (especially in patients who remain symptomatic after an adequate trial of artificial tear solutions)</li> <li>Exposure keratitis</li> <li>Decreased corneal sensitivity</li> <li>Recurrent corneal erosions</li> </ul>	Availability of an additional option for the treatment of dry eye.  The preferred option on the Advanced Control Formulary is Xiidra (lifitegrast).
<b>Maxidex</b> (dexamethasone) ophthalmic suspension	Topical/ Ophthalmic/ Anti-Inflammatories/ Steroidal	Maxidex is indicated for steroid responsive inflammatory conditions of the palpebral and bulbar conjunctiva, cornea, and anterior	Availability of additional ophthalmic anti-inflammatory options.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		segment of the globe such as allergic conjunctivitis, acne rosacea, superficial punctate keratitis, herpes zoster keratitis, iritis, cyclitis, selected infective conjunctivides when the inherent hazard of steroid use is accepted to obtain an advisable diminution in edema and inflammation; corneal injury from chemical, radiation, or thermal burns, or penetration of foreign bodies.	Preferred options on the Advanced Control Formulary include dexamethasone, loteprednol, prednisolone acetate 1%, Flarex (fluorometholone acetate), and FML S.O.P. (fluorometholone).
<b>Mirvaso</b> (brimonidine) topical gel	Topical/ Dermatology/ Rosacea	Mirvaso is indicated for the topical treatment of persistent (nontransient) facial erythema of rosacea in adults 18 years of age or older.	Availability of additional topical options for the treatment of rosacea.  Preferred option on the Advanced Control Formulary include azelaic acid gel, metronidazole, and Finacea Foam (azelaic acid).
<b>Neulasta, Neulasta Onpro</b> (pegfilgrastim) subcutaneous solution for injection	Hematologic/ Hematopoietic Growth Factors	Neulasta and Neulasta Onpro are indicated to: <ul style="list-style-type: none"> <li>• Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia</li> <li>• Increase survival in patients acutely exposed to myelosuppressive doses of</li> </ul>	Availability of an additional long-acting colony-stimulating factor option for those who are receiving myelosuppressive anti-cancer therapy.  The preferred option on the Advanced Control Formulary is Ziextenzo (pegfilgrastim-bmez).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		radiation (Hematopoietic Subsyndrome of Acute Radiation Syndrome)	
<b>NuvaRing</b> (ethinyl estradiol-etonogestrel) vaginal ring	Endocrine and Metabolic/ Contraceptives/ Vaginal	NuvaRing is indicated for use by women to prevent pregnancy.	Availability of additional vaginal contraceptive options.  Preferred options on the Advanced Control Formulary include ethinyl estradiol-etonogestrel and Annovera (segesterone acetate-ethinyl estradiol).
<b>Osphena</b> (ospemifene) oral tablet	Endocrine and Metabolic/ Menopausal Symptom Agents/ Oral	Osphena is indicated for: <ul style="list-style-type: none"> <li>• Treatment of moderate to severe dyspareunia, a symptom of vulvar and vaginal atrophy, due to menopause</li> <li>• Treatment of moderate to severe vaginal dryness, a symptom of vulvar and vaginal atrophy, due to menopause</li> </ul>	Availability of a generic oral estrogen option.  The preferred option on the Advanced Control Formulary is estradiol.
<b>Premarin</b> (estrogens, conjugated) oral tablet	Endocrine and Metabolic/ Menopausal Symptom Agents/ Oral	Premarin is indicated for: <ul style="list-style-type: none"> <li>• Treatment of moderate to severe vasomotor symptoms due to menopause</li> <li>• Treatment of moderate to severe vulvar and vaginal atrophy due to menopause</li> <li>• Treatment of hypoestrogenism due to hypogonadism, castration or primary ovarian failure</li> <li>• Treatment of breast cancer (for palliation inly) in appropriately-selected women and men with metastatic disease</li> </ul>	Availability of a generic oral estrogen option.  The preferred option on the Advanced Control Formulary is estradiol.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<ul style="list-style-type: none"> <li>Treatment of advanced androgen-dependent carcinoma of the prostate (for palliation only)</li> <li>Prevention of postmenopausal osteoporosis</li> </ul>	
<b>Premarin Cream</b> (estrogens, conjugated) vaginal cream	Endocrine and Metabolic/ Menopausal Symptom Agents/ Vaginal	Premarin Cream is indicated for: <ul style="list-style-type: none"> <li>Treatment of atrophic vaginitis and kraurosis vulvae</li> <li>Treatment of moderate to severe dyspareunia, a symptom of vulvar and vaginal atrophy, due to menopause</li> </ul>	Availability of additional vaginal estrogen options for the treatment of dyspareunia.  Preferred options on the Advanced Control Formulary include estradiol and Imvexxy (estradiol).
<b>Premphase</b> (estrogens, conjugated-medroxyprogesterone acetate) oral tablet	Endocrine and Metabolic/ Menopausal Symptom Agents/ Oral	Premphase is indicated in a woman with a uterus for: <ul style="list-style-type: none"> <li>Treatment of moderate to severe vasomotor symptoms due to menopause</li> <li>Treatment of moderate to severe vulvar and vaginal atrophy due to menopause</li> <li>Prevention of postmenopausal osteoporosis</li> </ul>	Availability of additional oral estrogen/progestin options.  Preferred options on the Advanced Control Formulary include estradiol-norethindrone and Bijuva (estradiol-progesterone).
<b>Prempro</b> (estrogens, conjugated-medroxyprogesterone acetate) oral tablet	Endocrine and Metabolic/ Menopausal Symptom Agents/ Oral	Prempro is indicated in a woman with a uterus for: <ul style="list-style-type: none"> <li>Treatment of moderate to severe vasomotor symptoms due to menopause</li> <li>Treatment of moderate to severe vulvar and vaginal atrophy due to menopause</li> </ul>	Availability of additional oral estrogen/progestin options.  Preferred options on the Advanced Control Formulary include estradiol-norethindrone and Bijuva (estradiol-progesterone).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		<ul style="list-style-type: none"> <li>Prevention of postmenopausal osteoporosis</li> </ul>	
<b>Prolensa</b> (bromfenac) ophthalmic solution	Topical/ Ophthalmic/ Anti-Inflammatories/ Nonsteroidal	Prolensa is indicated for the treatment of postoperative inflammation and reduction of ocular pain in patients who have undergone cataract surgery.	<p>Availability of generic ophthalmic anti-inflammatory options.</p> <p>Preferred options on the Advanced Control Formulary include bromfenac, diclofenac, and ketorolac.</p>
<b>Provad</b> (bacillus subtilis) oral capsule	Gastrointestinal/ Probiotics	Provad is a probiotic formulation for the clinical dietary management of suboptimal nutritional status in patients where advanced supplementation is required and nutritional supplementation in physiologically stressful conditions for maintenance of good health is needed.	<p>Availability of additional probiotic supplementation options.</p> <p>Consult doctor for preferred options on the Advanced Control Formulary.</p>
<b>Relistor</b> (methylnaltrexone) oral tablet	Gastrointestinal/ Opioid- Induced Constipation	Relistor tablet is indicated for the treatment of opioid-induced constipation (OIC) in adults with chronic non-cancer pain, including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation.	<p>Availability of additional options for the treatment of opioid-induced constipation.</p> <p>Preferred options on the Advanced Control Formulary include Movantik (naloxegol) and Symproic (naldemedine).</p>
<b>Relistor</b> (methylnaltrexone) subcutaneous solution for injection	Gastrointestinal/ Opioid- Induced Constipation	Relistor injection is indicated for <ul style="list-style-type: none"> <li>Treatment of opioid-induced constipation (OIC) in adults with chronic non-cancer pain, including patients with chronic pain related to prior cancer or its treatment who do not require</li> </ul>	<p>Availability of additional options for the treatment of opioid-induced constipation.</p> <p>Preferred options on the Advanced Control Formulary include Movantik (naloxegol) and Symproic (naldemedine).</p>

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		frequent (e.g., weekly) opioid dosage escalation <ul style="list-style-type: none"> <li>Treatment of OIC in adults with advanced illness or pain caused by active cancer who require opioid dosage escalation for palliative care</li> </ul>	
<b>Rexulti</b> (brexpiprazole) oral tablet	Central Nervous System/ Antipsychotics/ Atypicals	Rexulti is indicated for: <ul style="list-style-type: none"> <li>Use as an adjunctive therapy to antidepressants for the treatment of major depressive disorder (MDD)</li> <li>Treatment of schizophrenia</li> </ul>	Availability of additional oral antipsychotic options.  Preferred options on the Advanced Control Formulary include aripiprazole, clozapine, olanzapine, quetiapine, quetiapine ext-rel, risperidone, ziprasidone, Latuda (lurasidone), Saphris (asenapine), and Vraylar (cariprazine).
<b>Rhofade</b> (oxymetazoline) topical cream	Topical/ Dermatology/ Rosacea	Rhofade is indicated for the topical treatment of persistent facial erythema associated with rosacea in adults.	Availability of additional topical options for the treatment of rosacea.  Preferred options on the Advanced Control Formulary include azelaic acid gel, metronidazole, and Finacea Foam (azelaic acid).
<b>Risperdal CONSTA</b> (risperidone microspheres) intramuscular extended-release suspension for injection	Central Nervous System/ Antipsychotics/ Atypicals	Risperdal CONSTA is indicated: <ul style="list-style-type: none"> <li>For the treatment of schizophrenia</li> <li>As monotherapy or as adjunctive therapy to lithium or valproate for the maintenance treatment of Bipolar I Disorder</li> </ul>	Availability of additional long-acting injectable antipsychotic options.  Preferred options on the Advanced Control Formulary include Abilify Maintena (aripiprazole ext-rel injection) and Perseris (risperidone ext-rel injection).
<b>Signifor LAR</b> (pasireotide)	Endocrine and Metabolic/ Acromegaly	Signifor LAR is indicated for the treatment of:	Availability of an additional option for the treatment of acromegaly.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.



## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
intramuscular suspension for injection		<ul style="list-style-type: none"> <li>Patients with acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option</li> <li>Patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative</li> </ul>	The preferred option on the Advanced Control Formulary is Somatuline Depot (lanreotide acetate).
<b>Somavert</b> (pegvisomant) subcutaneous solution for injection	Endocrine and Metabolic/ Acromegaly	Somavert is indicated for the treatment of acromegaly in patients who have had an inadequate response to surgery or radiation therapy, or for whom these therapies are not appropriate.	<p>Availability of an additional option for the treatment of acromegaly.</p> <p>The preferred option on the Advanced Control Formulary is Somatuline Depot (lanreotide acetate).</p>
<b>Soolantra</b> (ivermectin) topical cream	Topical/ Dermatology/ Rosacea	Soolantra is indicated for the treatment of inflammatory lesions of rosacea.	<p>Availability of additional topical options for the treatment of rosacea.</p> <p>Preferred options on the Advanced Control Formulary include azelaic acid gel, metronidazole, and Finacea Foam (azelaic acid).</p>
<b>Suprep</b> (sodium sulfate-potassium sulfate-magnesium sulfate) oral solution	Gastrointestinal/ Laxatives	Suprep is indicated for cleansing of the colon in preparation for colonoscopy in adult and pediatric patients 12 years of age and older.	<p>Availability of additional options for colon cleansing prior to a colonoscopy.</p> <p>Preferred options on the Advanced Control Formulary include peg 3350-electrolytes and Clenpiq (sodium picosulfate-magnesium oxide-citric acid).</p>
<b>Tazorac</b> (tazarotene) topical cream	Topical/ Dermatology/ Acne/ Topical	<p>Tazorac cream 0.05% and 0.1% are indicated for:</p> <ul style="list-style-type: none"> <li>Topical treatment of plaque psoriasis</li> </ul>	Availability of additional options for the topical treatment of acne.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		Tazorac cream 0.1% is indicated for: <ul style="list-style-type: none"> <li>• Topical treatment of acne vulgaris</li> </ul>	<p>Preferred options on the Advanced Control Formulary include adapalene, benzoyl peroxide, clindamycin gel (except NDC 68682046275), clindamycin solution, clindamycin-benzoyl peroxide, erythromycin solution, erythromycin-benzoyl peroxide, tretinoin, Epiduo (adapalene-benzoyl peroxide), and Onexton (clindamycin-benzoyl peroxide).</p> <p>Availability of generic options for the topical treatment of plaque psoriasis.</p> <p>Preferred options on the Advanced Control Formulary include calcipotriene ointment and calcipotriene solution.</p>
<b>Tazorac</b> (tazarotene) topical gel	Topical/ Dermatology/ Acne/ Topical	<p>Tazorac gel 0.05% and 0.1% are indicated for:</p> <ul style="list-style-type: none"> <li>• Topical treatment of plaque psoriasis of up to 20% body surface area involvement</li> </ul> <p>Tazorac gel 0.1% is indicated for:</p> <ul style="list-style-type: none"> <li>• Topical treatment of mild to moderate facial acne vulgaris</li> </ul>	<p>Availability of additional options for the topical treatment of acne.</p> <p>Preferred options on the Advanced Control Formulary include adapalene, benzoyl peroxide, clindamycin gel (except NDC 68682046275), clindamycin solution, clindamycin-benzoyl peroxide, erythromycin solution, erythromycin-benzoyl peroxide, tretinoin, Epiduo (adapalene-benzoyl peroxide), and Onexton (clindamycin-benzoyl peroxide).</p> <p>Availability of generic options for the topical treatment of plaque psoriasis.</p>

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
			Preferred options on the Advanced Control Formulary include calcipotriene ointment and calcipotriene solution.
<b>Tecfidera</b> (dimethyl fumarate delayed-rel) oral delayed-release capsule	Central Nervous System/ Multiple Sclerosis Agents	Tecfidera is indicated for the treatment of relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.	<p>Availability of additional options for the treatment of relapsing forms of multiple sclerosis.</p> <p>Preferred options on the Advanced Control Formulary include dimethyl fumarate delayed-rel, glatiramer, Aubagio (teriflunomide), Betaseron (interferon beta-1b), Copaxone (glatiramer), Gilenya (fingolimod), Kesimpta (ofatumumab), Mayzent (siponimod), Ocrevus (ocrelizumab), Rebif (interferon beta-1a), Tysabri (natalizumab), Vumerity (diroximel fumarate delayed-rel), and Zeposia (ozanimod).</p>
<b>TobraDex</b> (tobramycin-dexamethasone) ophthalmic ointment, ophthalmic suspension	Topical/ Ophthalmic/ Anti-Infective / Anti-Inflammatory Combinations	TobraDex is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where superficial bacterial ocular infection or a risk of bacterial ocular infection exists.	<p>Availability of a generic ophthalmic anti-infective and anti-inflammatory option.</p> <p>The preferred option on the Advanced Control Formulary is tobramycin-dexamethasone.</p>
<b>TobraDex ST</b> (tobramycin-dexamethasone) ophthalmic suspension	Topical/ Ophthalmic/ Anti-Infective / Anti-Inflammatory Combinations	TobraDex ST is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where superficial bacterial ocular infection or a risk of bacterial ocular infection exists.	<p>Availability of a generic ophthalmic anti-infective and anti-inflammatory option.</p> <p>The preferred option on the Advanced Control Formulary is tobramycin-dexamethasone.</p>
<b>Tracleer</b> (bosentan) oral tablet, oral soluble tablet	Cardiovascular/ Pulmonary Arterial Hypertension/	Tracleer is indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization Group 1):	Availability of additional options for the treatment of pulmonary arterial hypertension.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
	Endothelin Receptor Antagonists	<ul style="list-style-type: none"> <li>• In adults to improve exercise ability and to decrease clinical worsening</li> <li>• In pediatric patients aged 3 years and older with idiopathic or congenital PAH to improve pulmonary vascular resistance, which is expected to result in an improvement in exercise ability</li> </ul>	Preferred options on the Advanced Control Formulary include ambrisentan, bosentan, and Opsumit (macitentan).
<b>Trulance</b> (plecanatide) oral tablet	Gastrointestinal/ Irritable Bowel Syndrome	Trulance is indicated in adults for the treatment of: <ul style="list-style-type: none"> <li>• Chronic idiopathic constipation (CIC)</li> <li>• Irritable bowel syndrome with constipation (IBS-C)</li> </ul>	Availability of an additional option for the treatment of chronic idiopathic constipation and irritable bowel syndrome with constipation.  The preferred option on the Advanced Control Formulary is Linzess (linaclotide).
<b>Udenyca</b> (pegfilgrastim-cbqv) subcutaneous solution for injection	Hematologic/ Hematopoietic Growth Factors	Udenyca is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.	Availability of an additional long-acting colony-stimulating factor option for those who are receiving myelosuppressive anti-cancer therapy.  The preferred option on the Advanced Control Formulary is Ziextenzo (pegfilgrastim-bmez).
<b>Varubi</b> (rolapitant) oral tablet	Gastrointestinal/ Antiemetics	Varubi is indicated in combination with other antiemetic agents in adults for the prevention of delayed nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy, including, but not limited to, highly emetogenic chemotherapy.	Availability of a generic option for the prevention of chemotherapy-induced nausea and vomiting.  The preferred option on the Advanced Control Formulary is aprepitant.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

## Advanced Control Formulary™ Change Summary Report Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Versacloz</b> (clozapine) oral suspension	Central Nervous System/ Antipsychotics/ Atypicals	Versacloz is indicated for: <ul style="list-style-type: none"> <li>• Treatment-resistant schizophrenia</li> <li>• Reducing suicidal behavior in patients with schizophrenia or schizoaffective disorder</li> </ul>	Availability of a generic oral antipsychotic option for treatment-resistant schizophrenia.  The preferred option on the Advanced Control Formulary is clozapine.
<b>Viibryd</b> (vilazodone) oral tablet, oral starter kit	Central Nervous System/ Antidepressants/ Selective Serotonin Reuptake Inhibitors (SSRIs)	Viibryd is indicated for the treatment of major depressive disorder (MDD) in adults.	Availability of additional selective serotonin reuptake inhibitor options.  Preferred options on the Advanced Control Formulary include citalopram, escitalopram, fluoxetine (except fluoxetine tablet 60 mg and generics for Sarafem), paroxetine HCl, paroxetine HCl ext-rel, sertraline, and Trintellix (vortioxetine).
<b>Visco-3</b> (sodium hyaluronate) intra-articular solution for injection	Analgesics/ Viscosupplements	Visco-3 is indicated for the treatment of pain in osteoarthritis of the knee in patients who have failed to respond adequately to conservative non-pharmacologic therapy and simple analgesics (e.g., acetaminophen).	Availability of additional viscosupplement options for osteoarthritis.  Preferred options on the Advanced Control Formulary include Durolane (sodium hyaluronate), Euflexxa (sodium hyaluronate), Gelsyn-3 (sodium hyaluronate), and Supartz FX (sodium hyaluronate).
<b>Zalvit</b> (multivitamin with iron) oral tablet	Nutritional/Supplements/ Vitamins and Minerals/ Prenatal Vitamins	Zalvit is used: <ul style="list-style-type: none"> <li>• Throughout pregnancy, during the postnatal period for both lactating and non-lactating mothers, and throughout the childbearing years</li> <li>• For improving the nutritional status of women prior to conception</li> </ul>	Availability of additional prenatal vitamin options.  Preferred options on the Advanced Control formulary include prenatal vitamins and Citranatal (prenatal vitamins with folic acid).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Zelac</b> (lactobacillus acidophilus and bifidobacterium animalis lactis) oral capsule	Gastrointestinal/ Probiotics	Zelac is a probiotic formulation for the clinical dietary management of suboptimal nutritional status in patients where advanced supplementation is required and nutritional supplementation in physiologically stressful conditions for maintenance of good health is needed.	Availability of additional probiotic supplementation options.  Consult doctor for preferred options on the Advanced Control Formulary.
<b>Zirgan</b> (ganciclovir) ophthalmic gel	Topical/ Ophthalmic/ Antivirals	Zirgan is indicated for the treatment of acute herpetic keratitis (dendritic ulcers).	Availability of a generic ophthalmic antiviral option.  The preferred option on the Advanced Control Formulary is trifluridine.
<b>Generic Agents:</b>			
<b>calcipotriene- betamethasone</b> topical ointment	Topical/ Dermatology/ Antipsoriatics	Calcipotriene-betamethasone ointment is indicated for the topical treatment of plaque psoriasis in patients 12 years of age and older.	Availability of additional topical options for the treatment of plaque psoriasis.  Preferred options on the Advanced Control Formulary include calcipotriene ointment or solution WITH desoximetasone, fluocinonide (except fluocinonide cream 0.1%), or Bryhali (halobetasol propionate); Duobrii (halobetasol propionate-tazarotene), Enstilar (calcipotriene-betamethasone dipropionate), or Taclonex (calcipotriene-betamethasone dipropionate).
<b>calcipotriene- betamethasone</b> topical suspension	Topical/ Dermatology/ Antipsoriatics	Calcipotriene-betamethasone suspension is indicated for the topical treatment of plaque psoriasis of the scalp in patients 12 years and older and plaque psoriasis of the scalp	Availability of additional topical options for the treatment of plaque psoriasis.  Preferred options on the Advanced Control Formulary include calcipotriene ointment or solution WITH

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.

Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
		and body in patients age 18 years and older.	desoximetasone, fluocinonide (except fluocinonide cream 0.1%), or Bryhali (halobetasol propionate); Duobrii (halobetasol propionate-tazarotene), Enstilar (calcipotriene-betamethasone dipropionate), or Taclonex (calcipotriene-betamethasone dipropionate).
<b>doxycycline monohydrate delayed-rel capsule</b> oral delayed-release capsule	Topical/ Dermatology/ Rosacea	Doxycycline monohydrate delayed-rel is indicated for the treatment of only inflammatory lesions (papules and pustules) of rosacea in adult patients.	Availability of an additional oral option for the treatment of rosacea.  The preferred option on the Advanced Control Formulary is Oracea (doxycycline monohydrate delayed-rel).
<b>isosorbide dinitrate 40 mg</b> oral tablet	Cardiovascular/ Nitrates/ Oral	Isosorbide dinitrate is indicated for the prevention of angina pectoris due to coronary artery disease.	Availability of additional generic oral nitrate options.  Preferred options on the Advanced Control Formulary include isosorbide dinitrate (except isosorbide dinitrate 40 mg) and isosorbide mononitrate.
<b>metaxalone 400 mg</b> oral tablet	Central Nervous System/ Musculoskeletal Therapy Agents	Metaxalone is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomforts associated with acute, painful musculoskeletal conditions.	Availability of additional generic options for the treatment of pain associated with acute musculoskeletal disorders.  Preferred options on the Advanced Control Formulary include carisoprodol, chlorzoxazone 500 mg (except NDC 73007001303), cyclobenzaprine (except cyclobenzaprine tablet 7.5 mg), metaxalone 800 mg, and methocarbamol (except NDCs 69036091010, 69036093090, 70868090190).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.



Advanced Control Formulary™  
 Change Summary Report  
 Effective 01-01-2021

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>methocarbamol (NDC 69036091010 only)</b> oral tablet	Central Nervous System/ Musculoskeletal Therapy Agents	Methocarbamol is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions.	Availability of additional generic options for the treatment of pain associated with acute musculoskeletal disorders.  Preferred options on the Advanced Control Formulary include carisoprodol, chlorzoxazone 500 mg (except NDC 73007001303), cyclobenzaprine (except cyclobenzaprine tablet 7.5 mg), metaxalone 800 mg, and methocarbamol (except NDCs 69036091010, 69036093090, 70868090190).
<b>oxymorphone ext-rel</b> oral extended-release tablet	Analgesics/ Opioid Analgesics	Oxymorphone ext-rel is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.	Availability of additional long-acting options for pain management.  Preferred options on the Advanced Control Formulary include fentanyl transdermal, hydrocodone ext-rel, methadone, morphine ext-rel, Nucynta ER (tapentadol ext-rel), and Xtampza ER (oxycodone ext-rel).

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.



**OTHER:**

Product	Therapeutic Category/ Subcategory	Indication	Options/Comments
<b>Tremfya</b> (guselkumab) subcutaneous solution for injection	Immunologic Agents/ Autoimmune Agents/ Psoriatic Arthritis	Tremfya is indicated for the treatment of adult patients with: <ul style="list-style-type: none"> <li>• Moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy</li> <li>• Active psoriatic arthritis</li> </ul>	Now excluded for psoriatic arthritis.  Preferred options on the Advanced Control Formulary include Cosentyx (secukinumab), Enbrel (etanercept), Humira (adalimumab), and Otezla (apremilast).  Note: Tremfya remains preferred for psoriasis.

This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers not affiliated with CVS Caremark. The information contained in this document is proprietary. The information may not be copied in whole or in part without written permission.  
 ©2020 CVS Caremark. All rights reserved.